



Hospital Quality Workgroup Starter Set Measures; Updated Feb. 21, 2007

| Quality Goal or Clinical Area                   | Indicators  | Forum Starter Set | HQA |
|---|---|-------------------|-----|
| <b>Acute Myocardial Infarction (AMI)</b>        | Aspirin at arrival  | √                 | √   |
|   | Aspirin prescribed at discharge   | √                 | √   |
|   | ACE-Inhibitor (ACE-1) or Angiotensin Receptor Blocker (ARBs) for left ventricular systolic dysfunction (LVSD) | √                 | √   |
|   | Beta blocker at arrival   | √                 | √   |
|   | Beta blocker prescribed at discharge  | √                 | √   |
|   | Thrombolytic agent received within 30 minutes of hospital arrival   | √                 | √   |
|   | Percutaneous Coronary Intervention received within 120 minutes of hospital arrival                            | √                 | √   |
|   | Adult smoking cessation advice/counseling   | √                 | √   |
| <b>Heart Failure (HF)</b>                       | Left ventricular function assessment  | √                 | √   |
|   | ACE-Inhibitor (ACE-1) or Angiotensin Receptor Blockers (ARB) for left ventricular systolic dysfunction (LVSD) | √                 | √   |
|   | Adult smoking cessation advice/counseling   | √                 | √   |
|   | Discharge instructions  | √                 | √   |
|   |   |                   |     |
| <b>Pneumonia</b>                                | Initial antibiotic received within 4 hrs of hospital arrival  | √                 | √   |
|   | Oxygenation assessment  | √                 | √   |
|   | Pneumococcal vaccination status   | √                 | √   |
|   | Blood cultures performed before first antibiotic received in hospital   | √                 | √   |
|   | Adult smoking cessation advice/counseling   | √                 | √   |
|   | Influenza vaccination status  | √                 | √   |
|   | Appropriate initial antibiotic selection  | √                 | √   |
| <b>Surgical Care Improvement Program (SCIP)</b> | Prophylactic antibiotic received within 1 hour prior to surgical incision                                     | √                 | √   |
|   | Prophylactic antibiotic selection for surgical patients   | √                 | √   |
|   | Prophylactic antibiotics discontinued within 24 hours after surgery end time                                  | √                 | √   |
|   | Surgery patients with recommended venous thromboembolism prophylaxis ordered (December)                       | √                 | √   |
|   |   |                   |     |